

Secondary non-Gaussianity and Cross-Correlation Analysis

Dipak Munshi^{1,2}, Patrick Valageas³, Asantha Cooray⁴ and Alan Heavens¹

¹Scottish Universities Physics Alliance (SUPA), Institute for Astronomy, University of Edinburgh, Blackford Hill, Edinburgh EH9 3HJ, UK

²School of Physics and Astronomy, Cardiff University, CF24 3AA

³Institut de Physique Théorique, CEA Saclay 91191, Gif-sur-Yvette, France

⁴Department of Physics and Astronomy, University of California, Irvine, CA 92697

18 July 2009, Revision: 0.9

ABSTRACT

We develop optimised estimators of two sorts of power spectra for fields defined on the sky, in the presence of partial sky coverage. The first is the cross-power spectrum of two fields on the sky; the second is the skew spectrum of three fields. The cross-power spectrum of the Cosmic Microwave Background (CMB) sky with tracers of large-scale-structure is useful as it provides valuable information on cosmological parameters. Numerous recent studies have proved the usefulness of cross-correlating CMB sky with external data sets, which probes the Integrated Sachs Wolfe Effect (ISW) at large angular scales and the Sunyaev Zeldovich (SZ) effect from hot gas in clusters at small angular scales. The skew spectrum, recently introduced by Munshi & Heavens (2009), is an optimised statistic which can be tuned to study a particular form of non-gaussianity, such as may arise in the early Universe, but which retains information on the nature of non-gaussianity. In this paper we develop the mathematical formalism for the skew spectrum of 3 different fields. When applied to the CMB, this allows us to explore the contamination of the skew spectrum by secondary sources of CMB fluctuations. Considering the three-point function, the study of the bispectrum provides valuable information regarding cross-correlation of secondaries with lensing of CMB with much higher significance compared to just the study involving CMB sky alone. After developing the analytical model we use them to study specific cases of cosmological interest which include cross-correlating CMB with various large scale tracers to probe ISW and SZ effects for cross spectral analysis. Next we use the formalism to study the signal-to-noise ratio for detection of the weak lensing of the CMB by cross-correlating it with different tracers as well as point sources for CMB experiments such as Planck.

Key words: : Cosmology– Cosmic microwave background– large-scale structure of Universe – Methods: analytical, statistical, numerical

INTRODUCTION

Observations of cosmic microwave background (CMB) and that of large scale structure carry complementary cosmological information. While all-sky CMB observations such as NASA’s WMAP¹ and ESA’s current Planck² experiments primarily probes the distribution of matter and radiation at redshift $z = 1300$, large-scale surveys tend to give us a window at lower redshift $z \sim 0$. The main advantage of cross-correlating such independent data sets lies in the fact that it is possible to highlight signals which may not be otherwise detected in individual data sets independently. Earlier studies in this direction include, Peiris & Spergel (2000) who carried out a detailed error forecast of such cross-correlation analysis for cosmological parameters. Clearly, for these tracers to be effective in constraining cosmology, they should be as numerous as possible to reduce the Poisson noise and the survey should cover as large a fraction of the sky as possible to reduce sample variance.

Various authors have used different external data sets with specific astrophysical tracers to trace the large-scale structure (LSS), with one of the main motivations being to detect the ISW effect as predicted for Λ CDM cosmology. Earlier studies in this direction include Fosabala & Gaztanaga (2004, 2006) who cross-correlated the SDSS Data Release 1 galaxies with the first-year full sky WMAP data. Nolta et al. (2003) cross-correlated the NVSS radio source catalogue with first-year full sky WMAP data. Scranton et al. (2003) who correlated Sloan Digital Survey against WMAP data. Boughn & Crittenden (2005, 2004a,b) used two tracers of the large-scale structure: the HEAO1 A2 full sky hard X-ray map and NVSS full sky radio galaxy survey. A maximum likelihood fit to both data sets yields a detection of an ISW amplitude at a level consistent with what is predicted by the Λ CDM cosmology. Most of these studies detected ISW effect at a level of 2-3 σ although the error analysis models and the statistic used were sometimes completely different (see Ho et. al. (2008) for tomographic studies involving ISW and Hirata et al. (2008) for weak lensing detection). The ISW effect remains one of the most direct and quantitative measure of the dark energy available to us today. Future all-sky missions such as Planck

¹ <http://map.gsfc.nasa.gov/>

² <http://www.rssd.esa.int/index.php?project=Planck>

will provide an excellent possibility to extend these studies to higher confidence regime. While the above studies are mainly focussed on large angular scales, where the ISW effect plays an important role, at small angular scales, the presence of clusters and probably the associated filamentary network in which they reside can also affect both CMB maps through the Sunyaev-Zeldovich effect (Sunyaev & Zeldovich 1980) as well as through the X-ray maps via bremsstrahlung. Cross-correlation analysis of the diffuse soft X-ray background maps of ROSAT with WMAP 1st year data were performed by (Diego, Silk, Silwa 2004; Diego, Silk, Silwa 2004). This study was motivated by the fact that hot gas in clusters can be more easily detected by cross-correlating X-ray and CMB maps. Although no evidence was found of this effect it opens the possibility of detecting such an effect in future high-resolution CMB maps. All these act as a motivation for development of a generic techniques to cross-correlate high-resolution CMB maps with other maps from LSS surveys. In this paper we focus on cross-correlating two or three different datasets, but the challenges are similar to those arising from a single dataset. For example, the estimation of the power spectrum from a single high-resolution map poses a formidable numerical problem in terms of computational requirements. Typically two different methods are followed. The first is the non-linear maximum likelihood method, or its quadratic variant, which can be applied to smoothed degraded maps, as it is not possible to directly invert a full pixel covariance matrix (Tegmark 1997). To circumvent this problem a pseudo- $\mathcal{C}_\ell$ (PCL) technique was invented (Hivon et al. 2002) which is unbiased though remains suboptimal. In recent analysis Efstathiou (2004, 2006) has shown how to optimize these estimators which can then be used to analyse high-resolution maps in a very fast and accurate way. We generalize the PCL-based approach here to compute the cross-correlation of different data sets. The method developed here is completely general and can be applied to an arbitrary number of data sets. For example, our formalism can analyse the degree of cross-correlation among various CMB surveys observing the same region of the sky with different noise levels and survey strategies.

For near-Gaussian fields, two-point analysis from any cosmological survey provides the bulk of the cosmological information. Nevertheless, going one step further, at the level of three-point correlation, the detection of departure from Gaussianity in the CMB can probe both primary non-Gaussianity see (e.g. Munshi & Heavens (2009)) as well as the mode-coupling effects due to secondaries. The possibility of further improving a detection of primordial non-Gaussianity with CMB maps, given current hints with WMAP data (Yadav & Wandelt 2008; Smith, Senatore & Zaldarriaga 2009), provides further motivation in this direction. One of the prominent contributions to the secondary non-Gaussianity is the coupling of weak lensing and sources of secondary contributions such as SZ (Goldberg & Spergel 1999; Cooray & Hu 2000). Although weak lensing produces a characteristic signature in the CMB angular power spectrum, its detection has proved to be difficult internally from CMB power spectrum alone. The non-Gaussianity imprinted by lensing into the primordial CMB remains below the detection level of current experiments, although with Planck the situation is likely to improve. The difficulty originates mainly due to the fact that such detections are linked to the four-point statistics of the lensing potential. However cross-correlating CMB data with external tracers means lensing signals can be probed at the level of the mixed bispectrum. After the first unsuccessful attempt to cross-correlate WMAP against SDSS, recent efforts by Smith, Zahn & Dore (2007) have found a clear signal of weak lensing of the CMB, by cross-correlating WMAP against NVSS which covers a significant fraction of the sky. Their work also underlines the link between three-point statistics estimators and the estimators for weak lensing effects on CMB.

The study of non-Gaussianity is primarily focused on the bispectrum (Heavens 1998), however in practice it is difficult to probe the entire configuration dependence in the harmonic space from noisy data. The cumulant correlators are multi-point correlators collapsed to probe two-point statistic. These were introduced in the context of analyzing galaxy clustering by Szapudi & Szalay (1999), and were later found to be useful for analyzing projected surveys such as APM (Munshi, Melott & Coles 2000). Being two-point statistics they can be analyzed in the multipole space by defining an associated power-spectrum. Recent studies by Cooray (2006) and Cooray, Li & Melchiorri (2008) have shown its wider applicability including e.g. in 21cm studies. However, the multi-spectrum elements defined in multipole space are difficult to estimate directly from the data because of their complicated response to partial sky coverage and inhomogeneous noise, as well as associated high redundancy in the information content. However such issues are well understood in the context of power-spectrum analysis. Borrowing from previous results, in this paper we show how the cross-power spectrum and the skew spectrum can be studied in real data in an optimal way. We concentrate on two effects: the cross-correlation power spectrum, which is recovered by cross-correlating two different (but possibly correlated) data sets, focussing on weak lensing effects on the CMB, and secondly the contributions to the skew spectrum from foreground effects. The relation of such cross-power spectrum estimators with higher-order multi-spectra such as the bispectrum is also discussed in the context of methods known as pseudo- $\mathcal{C}_\ell$ s and quadratic estimators. We derive the error-covariance matrices and discuss their validity in the signal- and noise-dominated regimes and comment on their relationship to the Fisher matrix.

This layout of the paper is as follows: in §2 we use the formalism based on Pseudo- $\mathcal{C}_\ell$ analysis for power-spectra to study the cross-correlation power spectrum of different data sets. While we keep the analysis completely general, it is specialized for the case of near all-sky analysis and use it to compute the signal-to-noise and the covariance of estimated $\mathcal{C}_\ell$ s for various tracers with Planck-type all-sky experiments. Possibilities of using various weights which can make the pseudo- $\mathcal{C}_\ell$ approach near optimal in limiting cases of the signal-dominated regime or the noise-dominated regime are also discussed. In §3 we continue our discussion on Pseudo- $\mathcal{C}_\ell$ but generalize it to the analysis of the skew spectrum. Such an estimator can handle the partial sky coverage and noise in a very straightforward way. However in general it remains sub-optimal. In the high- l regime where mode-mode coupling can be modelled using a fraction of sky f_{sky} proxy one can make such an estimator nearly optimum using a suitable weighting. After a very brief introduction to various physical effects in §4 which introduce mode-mode coupling that leads to CMB bispectra, we move on to develop a crude but fast estimator for the skew spectrum in §5. Section §6 is devoted to developing the mixed bispectrum analysis in an optimal way by introducing inverse covariance weighting of the data. We analyze both one-point and two-point collapsed bispectral analysis. The one-point estimator or the mixed skewness is introduced - being a one point estimator it compresses all available information in a bispectrum to a single number. Next, we introduce the mixed skew spectrum which compresses various components of a bispectrum to a power spectrum in an optimum way. Next, in §7, the general formalism of bispectral analysis is used for specific cases of interest.

2 THE PSEUDO- $\mathcal{C}_\ell$ ESTIMATOR FOR CROSS-CORRELATION ANALYSIS

In this section we generalise results from pseudo- $\mathcal{C}_\ell$ power spectrum estimation of a single field with partial sky coverage, to cross-power spectra of two fields. For two fields Φ^X, Φ^Y defined on the sky, the pseudo- $\mathcal{C}_\ell$ estimators are constructed from the spherical harmonic transforms $a_{lm}^{X,Y}$ over the partial sky, where the fields are assumed to take zero value in unobserved regions.




Figure 1. The power-spectra C_l s are plotted for various analysis. In the left panel we show the power spectrum corresponding to a LSS tracer such as NVSS and that of CMB ISW. The power spectra associated with the two is also depicted. In the right panel we show SZ and ISW power-spectrum and their cross-correlation. The CMB power spectrum is also plotted in both panels. SZ curves include only the part that is correlated with the large-scale density field.

The $\tilde{a}_{lm}^{X,Y}$ are related to the true all-sky spherical harmonics $a_{\ell m}^{X,Y}$ by a linear transformation, via the transformation matrix $K_{lm'l'm'}$, and where possible direct inversion to get a_{lm} is much faster than maximum-likelihood analysis. Using a suitable choice of weighting function for the data this estimation can also be made nearly optimal.

2.1 Estimator for $\tilde{C}_l^{X,Y}$

The transformation from the pixel-space to harmonic domain can be expressed as follows:

$$\tilde{a}_{lm}^{X,Y} = \sum_{pixels} \Phi^{X,Y}(\hat{\Omega}_i) w(\hat{\Omega}_i)^{X,Y} \Omega_p(\hat{\Omega}_i)^{X,Y} Y_{lm}(\hat{\Omega}_i) = \sum_{l'm'} a_{l'm'}^{X,Y} K_{lm'l'm'}^{X,Y}. \quad (1)$$

Here $w(\hat{\Omega}_i)$ denotes the pixel space weight, $Y_{lm}(\hat{\Omega}_i)$ represents the spherical harmonic and $\Omega_p(\hat{\Omega}_i)$ is the pixel area (which we will assume independent of the pixel position). Expanding the weighting function in a spherical basis one can write down the coupling matrix $K_{l_1 m_1 l_2 m_2}$ which encodes all information regarding mode-mode coupling due to partial sky coverage as (see e.g. Hivon et al. (2002) for detailed derivation):

$$\begin{aligned} K_{l_1 m_1 l_2 m_2} &= \int w(\hat{\Omega}) Y_{l_1 m_1}(\hat{\Omega}) Y_{l_2 m_2}(\hat{\Omega}) d\hat{\Omega} \\ &= \sum_{l_3 m_3} \tilde{w}_{l_3 m_3} \left(\frac{(2l_1 + 1)(2l_2 + 1)(2l_3 + 1)}{4\pi} \right)^{1/2} \begin{pmatrix} l_1 & l_2 & l_3 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l_1 & l_2 & l_3 \\ m_1 & m_2 & m_3 \end{pmatrix}, \end{aligned} \quad (2)$$

where $\tilde{w}_{lm}$ is the transform of the window or (arbitrary) weighting function. The matrices represents $3J$ functions. The quantum numbers l and m need to satisfy certain conditions for the $3J$ functions to have non-vanishing values. The pixel area of maps Φ^X and Φ^Y will be denoted by and the associated weights with each pixels will be left arbitrary $w_i^{X,Y}$. Adopting the notation of Efstathiou (2004), we write the pseudo- $\mathcal{C}_\ell^{X,Y,XY}$ in terms of the underlying true power-spectrum $\mathcal{C}_\ell^{X,Y,XY}$.

Defining the following:

$$\langle \tilde{\mathcal{C}}_\ell^X \rangle = \frac{1}{(2\ell + 1)} \sum_m |\tilde{a}_{lm}^X|^2; \quad \langle \tilde{\mathcal{C}}_\ell^{XY} \rangle = \frac{1}{(2\ell + 1)} \sum_m \text{Real}(\tilde{a}_{lm}^X * \tilde{a}_{lm}^Y) \quad (3)$$

and similarly for $\langle \tilde{\mathcal{C}}_\ell^X \rangle$, we have

$$\langle \tilde{\mathcal{C}}_\ell^\alpha \rangle = \sum_{\ell'} C_{\ell'}^\alpha M_{\ell\ell'}^\alpha, \quad (4)$$

where $\alpha = X, Y$ or XY , and we can estimate the true covariance matrices as

$$\hat{\mathcal{C}}_\ell^\alpha = (M^{-1})_{\ell\ell'}^\alpha \tilde{\mathcal{C}}_{\ell'}^\alpha. \quad (5)$$

where the matrix M can be expressed in terms of $3J$ symbols as (Hivon et al. 2002):

$$M_{\ell_1 \ell_2}^\alpha = (2\ell_2 + 1) \sum_{\ell_3} \frac{(2\ell_3 + 1)}{4\pi} \tilde{W}_{l_3}^\alpha \begin{pmatrix} \ell_3 & \ell_2 & \ell_1 \\ 0 & 0 & 0 \end{pmatrix}^2. \quad (6)$$

and $W_l = \frac{1}{2l+1} \sum_m |w_{lm}|^2$ is the power spectra associated with the mask. Note that the transformation matrices $M^{X,Y}$ depends on the power-spectrum of the weighting function $w_{lm}^{X,Y}$, whereas the matrix M^{XY} for cross-power spectra is determined by the cross-power spectra $\tilde{W}_{l_3}^{XY}$ of two weighting functions. Independent of the choice of weights the estimators $\mathcal{C}_\ell^\alpha$ remain unbiased.




Figure 2. The variance for estimated C_l s are plotted. The variance for estimated C_l corresponding to ISW, SZ and their cross-correlation are plotted in the left panel. The ISW vs local tracers (NVSS type) analysis is plotted in the right. The number density of galaxies for NVSS-type survey was taken to be $\bar{N} = 7 \times 10^8$ per Steradian. For CMB a Planck type experiment was assumed. Results are for all-sky surveys. Results plotted are for $f_{sky} = 1$. For near all sky survey, the variances will scale linearly with f_{sky} .

2.2 Covariances of Pseudo- C_ℓ s

The pseudo- C_ℓ s are unbiased. The variances of these estimators can be computed analytically for arbitrary sky coverage and a non-uniform Gaussian noise distribution. The deviation of the estimated $\tilde{C}_l$ from the ensemble average $\langle \tilde{C}_l \rangle$ is denoted by $\delta \tilde{C}_l$.

$$\delta \tilde{C}_l^\alpha = \tilde{C}_l^\alpha - \langle \tilde{C}_l^\alpha \rangle; \quad \alpha \in X, Y, XY \quad (7)$$

We are concerned here with the computation of the covariance of estimated C_l s. We begin by defining the covariance matrix:

$$\langle \delta \tilde{C}_l^\alpha \delta \tilde{C}_{l'}^\beta \rangle = \langle \tilde{C}_l^\alpha \tilde{C}_{l'}^\beta \rangle - \langle \tilde{C}_l^\alpha \rangle \langle \tilde{C}_{l'}^\beta \rangle; \quad \alpha, \beta \in X, Y, XY \quad (8)$$

The covariance of C_l s from individual surveys Φ^X or Φ^Y can be expressed as follows (Efstathiou 2004).

$$\begin{aligned} \langle \delta \tilde{C}_\ell^X \delta \tilde{C}_{\ell'}^Y \rangle = & \sum_L \left\{ C_\ell^X C_{\ell'}^Y \frac{1}{2L+1} \sum_M |w_{LM}^X|^2 |w_{LM}^Y|^2 + C_\ell^X \frac{1}{2L+1} \sum_M |w_{LM}^X|^2 |(w\sigma^2)_{LM}^Y|^2 \right. \\ & \left. + \frac{1}{2L+1} \sum_M |w_{LM}^X|^2 |(w\sigma^2)_{LM}^Y|^2 \right\} \begin{pmatrix} L & \ell & \ell' \\ 0 & 0 & 0 \end{pmatrix}^2. \end{aligned} \quad (9)$$

Extending the above results similarly the covariance of C_l^X for the cross-power-spectrum can be expressed as:

$$\langle \delta \tilde{C}_\ell^{XY} \delta \tilde{C}_{\ell'}^{XY} \rangle = \sum_L \left\{ C_\ell^X C_{\ell'}^Y \frac{1}{2L+1} \sum_M |w_{LM}^X|^2 |w_{LM}^Y|^2 + C_\ell^X \frac{1}{2L+1} \sum_M |w_{LM}^X|^2 |(w^2\sigma^2)_{LM}^X|^2 \right. \quad (10)$$

$$\left. + C_{\ell'}^Y \frac{1}{2L+1} \sum_M |w_{LM}^Y|^2 |(w^2\sigma^2)_{LM}^Y|^2 + \frac{1}{2L+1} \sum_M |(w^2\sigma^2)_{LM}^X|^2 |(w^2\sigma^2)_{LM}^Y|^2 \right\} \begin{pmatrix} L & \ell & \ell' \\ 0 & 0 & 0 \end{pmatrix}^2, \quad (11)$$

In our derivation we have assumed that all three power spectra are being estimated from the data simultaneously.

The three off-diagonal terms can be similarly expressed as:

$$\langle \delta \tilde{C}_\ell^X \delta \tilde{C}_{\ell'}^Y \rangle = \sum_L (C_\ell^X)^2 \frac{1}{2L+1} |w_{LM}^X w_{LM}^{Y*}|^2 \left(\frac{2L+1}{2\pi} \right) \begin{pmatrix} L & \ell & \ell' \\ 0 & 0 & 0 \end{pmatrix}^2, \quad (12)$$

$$\langle \delta \tilde{C}_\ell^{XY} \delta \tilde{C}_{\ell'}^Y \rangle = \sum_L C_\ell^{XY} C_{\ell'}^Y \left\{ \frac{1}{2L+1} \sum_M |w_{LM}^X w_{LM}^{Y*}| |w_{LM}^X|^2 + C_\ell^{XY} \frac{1}{2L+1} \sum_M |(w^2\sigma^2)_{LM}^X w_{LM}^{Y*}|^2 \right\} \left(\frac{2L+1}{2\pi} \right) \begin{pmatrix} L & \ell & \ell' \\ 0 & 0 & 0 \end{pmatrix}^2. \quad (13)$$

Here we have introduced following notations:

$$w_{lm}^X = \int d\Omega w^X(\hat{\Omega}) Y_{lm}(\hat{\Omega}) \quad (14)$$

$$(w\sigma^2)_{lm}^X = \int d\hat{\Omega} w^X(\hat{\Omega}) \sigma_X^2(\Omega) Y_{lm}(\hat{\Omega}) \quad (15)$$

$$(w^2\sigma^2)_{lm}^X = \int d\hat{\Omega} w^X(\hat{\Omega})^2 \sigma_X^2(\hat{\Omega}) Y_{lm}(\hat{\Omega}). \quad (16)$$




Figure 3. The cumulative signal to noise for ISW cross LSS (right panel) and SZ cross ISW (left panel) are plotted as a function of l . The results are obtained by using the covariances of C_l s plotted in previous plot. The S/N will degrade linearly with f_{sky} . Plots correspond to $f_{sky} = 1$.

Similar expression hold for the second survey and the cross terms for product of two surveys are also needed to derive the error covariance matrices. Finally the error covariances associated with deconvolved estimators $\hat{C}_l$ can be expressed in terms of that of the convolved estimators $\tilde{C}_l$ as follows:

$$\begin{pmatrix} \langle \delta \hat{C}_L^X \delta \hat{C}_{L'}^X \rangle & \langle \delta \hat{C}_L^X \delta \hat{C}_{L'}^Y \rangle \\ \langle \delta \hat{C}_L^Y \delta \hat{C}_{L'}^X \rangle & \langle \delta \hat{C}_L^Y \delta \hat{C}_{L'}^Y \rangle \end{pmatrix} = \begin{pmatrix} M_{LL}^{XX} & M_{LL}^{XY} \\ M_{LL}^{YX} & M_{LL}^{YY} \end{pmatrix}^{-1} \begin{pmatrix} \langle \delta \tilde{C}_l^X \delta \tilde{C}_{l'}^X \rangle & \langle \delta \tilde{C}_l^X \delta \tilde{C}_{l'}^Y \rangle \\ \langle \delta \tilde{C}_l^Y \delta \tilde{C}_{l'}^X \rangle & \langle \delta \tilde{C}_l^Y \delta \tilde{C}_{l'}^Y \rangle \end{pmatrix} \begin{pmatrix} M_{L'L'}^{XX} & M_{L'L'}^{XY} \\ M_{L'L'}^{YX} & M_{L'L'}^{YY} \end{pmatrix}^{-1T} \quad (17)$$

In deriving these results it is assumed that the coverage of the sky is near complete. This will mean that the windows associated with the various couplings are sharper than any features in the power spectra. The shape of the mask and the noise covariance properties are quite general at this stage. If the C_l s of individual data sets are known from independent estimations then cross spectra deconvolution of the cross-spectra $C_l^{X,X}$ can be simply written as:

$$\langle \hat{C}_L^{X,Y} \hat{C}_{L'}^{X,Y} \rangle = \sum_{l'l'} [M^{X,Y}]_{Ll}^{-1} \langle \tilde{C}_l^{X,Y} \tilde{C}_{l'}^{X,Y} \rangle [M^Y]_{L'l'}^{-1}. \quad (18)$$

In the limiting situation when the survey area covers almost the entire sky these equation takes a much simpler form which are in common use in the literature. If f_{sky} is the fraction of the sky covered then one can write:

$$\langle \delta C_l^{XY} \delta C_{l'}^{XY} \rangle = f_{sky} \frac{1}{(2l+1)} [C_l^X C_{l'}^Y + (C_l^{XY})^2] \delta_{ll'}; \quad \langle \delta C_l^{XY} \delta C_{l'}^{X,Y} \rangle = f_{sky} \frac{2}{(2l+1)} (C_l^{X,Y} C_{l'}^{XY}) \delta_{ll'}. \quad (19)$$

The C_l s in these expression are the total $C_l = C_l^S + C_l^N$ which takes contribution from both signal and noise C_l s.

3 THE PSEUDO- C_ℓ ESTIMATOR FOR MIXED BISPECTRUM ANALYSIS

The statistics of temperature fluctuations in the sky are very nearly Gaussian, but small departures from Gaussianity can put constraints on early universe scenarios. Secondary non-Gaussianity on the other hand can provide valuable information to distinguish structure formation scenarios, and when used with constraints from the power spectrum it can be a very valuable tool. However estimation of the bispectrum for each triplet of harmonics modes can be difficult to perform numerically. Munshi & Heavens (2009) introduced a limited data compression method for 3-point functions which reduces the data to a single function (the skew spectrum), and which can be made optimal for estimating a bispectrum form. In the same spirit, we define a pseudo-skew spectrum for three arbitrary fields defined on a cut sky, and show how it is related to the skew spectrum on the uncut sky. The PCL-based approach described here is not optimal, however, it can be made optimal with suitable choice of weights.

Let us assume that we have three fields which are defined over the observed sky. The product of two of these fields as $X(\hat{\Omega})Y(\hat{\Omega})$ has an associated mask which, we denote as $w_A(\Omega)$, and which is a product of two masks associated with the individual fields. Analogously, the third field $Z(\Omega)$ is observed with a mask $w_B(\Omega)$.

From the harmonic transforms of these fields we study the skew spectrum and express it in terms of the mixed bispectra of fields X, Y and Z, $B_{l_1 l_2 l_3}^{XYZ}$. We develop this generally, but the results we derive will be useful for the study of primordial non-Gaussianity. Here we consider a single field (the CMB), but it is a field with contributions from various components, and the skew spectrum contains terms from various triplet of different (or repeated) fields.

3.1 Estimator for $\tilde{C}_l^{XY,Z}$

3.1.1 All-Sky Analysis

We start by introducing the power spectrum $\tilde{C}_l^{XY,Z}$ associated with the cross-correlation of the product map $X(\hat{\Omega})Y(\hat{\Omega})$ and $Z(\hat{\Omega})$. In the absence of sky-cuts and instrumental noise we can write:

6 Munshi, Valageas, Cooray & Heavens

$$\hat{C}_l^{XY,Z} = \frac{1}{2l+1} \sum_m a_{lm}^{XY} a_{lm}^{Z*} \quad (20)$$

where a_{lm}^{XY} is the spherical harmonic transform of XY .

Assuming homogeneity and isotropy, the correlation function $\mathcal{C}(\Omega, \Omega')$ of $X(\Omega)Y(\Omega)$ and $Z(\Omega)$ can be written in terms of $\hat{C}_l^{XY,Z}$

$$\mathcal{C}(\Omega, \Omega') \equiv \langle X(\Omega)Y(\Omega)Z(\Omega') \rangle = \sum_{l_1 m_1, l_2 m_2} \langle a_{l_1 m_1}^{XY} a_{l_2 m_2}^Z \rangle Y_{l_1 m_1}(\Omega) Y_{l_2 m_2}(\Omega') = \frac{1}{4\pi} \sum_l (2l+1) P_l(\cos(\hat{\Omega} \cdot \hat{\Omega}')) \hat{C}_l^{XY,Z}. \quad (21)$$

Here $P_l(\hat{\Omega} \cdot \hat{\Omega}')$ is a Legendre polynomial of order l . The three-point correlation function in the harmonic domain can similarly be used to introduce the mixed bispectrum $B_{l_1 l_2 l_3}^{XYZ}$ for the related fields. Assuming statistical isotropy,

$$\langle a_{l_1 m_1}^X a_{l_2 m_2}^Y a_{l_3 m_3}^Z \rangle = \begin{pmatrix} l_1 & l_2 & l_3 \\ m_1 & m_2 & m_3 \end{pmatrix} B_{l_1 l_2 l_3}^{XYZ}. \quad (22)$$

Our aim is to compute the cross-correlation power spectra of the product field $X(\Omega)Y(\Omega)$ and $Z(\Omega)$. Using a harmonic decomposition we can relate the multipoles a_{lm}^{XY} with multipoles $a_{l'm'}^X$ and $a_{l''m''}^Y$:

$$\begin{aligned} a_{lm}^{XY} &= \int d\hat{\Omega} Y_{lm}^*(\hat{\Omega}) X(\hat{\Omega}) Y(\hat{\Omega}) = \sum_{l'm'} \sum_{l''m''} a_{l'm'}^X a_{l''m''}^Y \int d\hat{\Omega} Y_{lm}^*(\hat{\Omega}) Y_{l'm'}(\hat{\Omega}) Y_{l''m''}(\hat{\Omega}) \\ &= \sum_{l'm'} \sum_{l''m''} a_{l'm'}^X a_{l''m''}^Y \sqrt{\frac{(2l_1+1)(2l_2+1)(2l_3+1)}{4\pi}} \begin{pmatrix} l & l' & l'' \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l & l' & l'' \\ m & m' & m'' \end{pmatrix}. \end{aligned} \quad (23)$$

Contracting with the multipole of the remaining field $a_{l'm'}^Z$ we can see that it directly probes the mixed bispectrum associated with these three different fields (Cooray 2001a).

$$\langle a_{lm}^{XY} a_{l'm'}^{Z*} \rangle \equiv C_l^{XY,Z} \delta_{ll'} \delta_{mm'}; \quad C_l^{XY,Z} = \sum_{l_1, l_2} B_{l l_1 l_2}^{XYZ} \sqrt{\frac{(2l_1+1)(2l_2+1)}{4\pi(2l+1)}} \begin{pmatrix} l_1 & l_2 & l \\ 0 & 0 & 0 \end{pmatrix} \quad (24)$$

Since the bispectrum is determined by triangular configuration in the multipole space (l_1, l_2, l) , the power spectrum C_l defined above captures information about all possible triangular configuration when one of its sides is fixed at length l . However this data compression is not done optimally as it does not weight the contributions from each bispectrum components with their inverse variance. The error covariance matrix can be computed exactly and depends on higher-order moments of signal and noise, as well as their cross-correlations.

3.1.2 Partial sky Coverage

It is possible to extend the above result to take into account partial sky coverage. Assuming the composite map $X(\hat{\Omega})Y(\hat{\Omega})$ is masked with arbitrary mask $w_A(\hat{\Omega})$ and the map $Z(\hat{\Omega})$ is masked with $w_B(\hat{\Omega})$ we can write the cut-sky multipoles $\tilde{a}_{lm}^{XY}$ and $\tilde{a}_{lm}^Z$ in terms of their all sky counterparts as well as the multipoles associated with the mask multipoles follows:

$$\tilde{a}_{lm}^{XY} = \int X(\hat{\Omega}) Y(\hat{\Omega}) w_A(\hat{\Omega}) Y_{lm}^*(\hat{\Omega}) d\hat{\Omega}; \quad \tilde{a}_{lm}^Z = \int Z(\hat{\Omega}) w_B(\hat{\Omega}) Y_{lm}^*(\hat{\Omega}) d\hat{\Omega} \quad (25)$$

$$\begin{aligned} \tilde{a}_{lm}^{XY} &= \sum_{l_1 m_1} \sum_{l_2 m_2} \sum_{l_A m_A} a_{l_1 m_1}^X a_{l_2 m_2}^Y w_{l_A m_A} \int d\hat{\Omega} Y_{l_1 m_1}(\hat{\Omega}) Y_{l_2 m_2}(\hat{\Omega}) Y_{l_A m_A}(\hat{\Omega}) \\ \tilde{a}_{lm}^Z &= \sum_{l_3 m_3} \sum_{l_B m_B} a_{l_3 m_3}^Z w_{l_B m_B} \int d\hat{\Omega} Y_{l_3 m_3}(\hat{\Omega}) Y_{l_B m_B}(\hat{\Omega}). \end{aligned} \quad (26)$$

Here $B_{l_1 l_2 l_3}$ is the angle-averaged bispectrum and the functions b_{l_i} represent the effects of pixellisation as well as beam smoothing. For partial sky coverage one can obtain after tedious but straight forward algebra:

$$\tilde{C}_l^{XY,Z} = \sum_{l'} \frac{2l'+1}{4\pi} \sum_{l''} \frac{(2l''+1)}{4\pi} \begin{pmatrix} l & l' & l'' \\ 0 & 0 & 0 \end{pmatrix}^2 |w_{l''}|^2 \sum_{l_1, l_2} B_{l l_1 l_2}^{XYZ} \sqrt{\frac{(2l_1+1)(2l_2+1)}{(2l'+1)4\pi}} \begin{pmatrix} l_1 & l_2 & l' \\ 0 & 0 & 0 \end{pmatrix} \equiv M_{ll'} \hat{C}_{l'}^{XY,Z}. \quad (27)$$

This is one of the important results in this paper. It shows how the pseudo-skew spectrum is related to the all-sky skew spectrum, and is a computationally-efficient way to estimate the latter. By suitable choice of weight functions it can be made optimal. It is valid for a completely general mask where w_l represents the spherical transform of the mask.

The transformation matrix $M_{ll'}$ used here is the same as that we introduced for the recovery of cross power spectra. The power spectra associated with the mask w_l , plays the same role in construction of $M_{ll'}$. For simplicity we have assumed that different data sets have the same mask but it

is trivial to generalise for two different masks. A detailed comparison of level of sub-optimality will be compared with numerical simulations in an accompanying paper. This analysis is complementary to work by Chen & Szapudi (2006) where a similar sub-optimal estimator was used to study non-Gaussianity.

4 CMB SECONDARY BISPECTRUM

The formalism developed so far is quite general and can handle mixed bispectra of different kinds. The main goal was to relate the skew spectrum with the corresponding mixed bispectrum. To make concrete predictions we need to consider a specific form for the bispectrum. Following Spergel & Goldberg (1999), Goldberg & Spergel (1999) and Cooray & Hu (2000) we expand the observed temperature anisotropy $\delta T(\hat{\Omega})$ in terms of the primary anisotropy δT_P , and due to lensing of primary, δT_L , and the other secondaries from coupling large-scale structure, δT_S .

$$\delta T(\hat{\Omega}) = \delta T_P(\hat{\Omega}) + \delta T_L(\hat{\Omega}) + \delta T_S(\hat{\Omega}). \quad (28)$$

Expanding the respective terms in spherical harmonics we can write:

$$\delta T_P(\hat{\Omega}) \equiv \sum_{lm} a_{lm} Y_{lm}(\hat{\Omega}); \quad \delta T_L(\hat{\Omega}) \equiv \sum_{lm} a_{lm} \nabla \Theta(\hat{\Omega}) \cdot \nabla T_S(\hat{\Omega}); \quad \delta T_S(\hat{\Omega}) \equiv \sum_{lm} b_{lm} Y_{lm}(\hat{\Omega}). \quad (29)$$

The harmonic coefficients b_{lm} are associated with the expansion of the secondary anisotropies $\delta T_S(\hat{\Omega})$. The secondary bispectrum for the CMB then takes contributions from many products of P, L, S terms. For example, one term arises from products of $\delta T_P \delta T_L \delta T_S$:

$$\begin{aligned} B_{l_1 l_2 l_3}^{PLS} &\equiv \sum_{m_1 m_2 m_3} \begin{pmatrix} l_1 & l_2 & l_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \int \langle \delta T_P(\hat{\Omega}_1) \delta T_L(\hat{\Omega}_2) \delta T_S(\hat{\Omega}_3) \rangle Y_{l_1 m_1}(\hat{\Omega}_1) Y_{l_2 m_2}(\hat{\Omega}_2) Y_{l_3 m_3}(\hat{\Omega}_3) d\hat{\Omega}_1 d\hat{\Omega}_2 d\hat{\Omega}_3 \\ &\equiv \sum_{m_1 m_2 m_3} \begin{pmatrix} l_1 & l_2 & l_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \langle (\delta T_P)_{l_1 m_1} (\delta T_L)_{l_2 m_2} (\delta T_S)_{l_3 m_3} \rangle. \end{aligned} \quad (30)$$

It is possible to invert the relation using isotropy of the background cosmology:

$$\langle (\delta T_P)_{l_1 m_1} (\delta T_P)_{l_2 m_2} (\delta T_P)_{l_3 m_3} \rangle = \begin{pmatrix} l_1 & l_2 & l_3 \\ m_1 & m_2 & m_3 \end{pmatrix} B_{l_1 l_2 l_3}. \quad (31)$$

Explicit calculations, detailed in Goldberg & Spergel (1999) and Cooray & Hu (2000), found the mixed bispectrum to be of the following form:

$$B_{l_1 l_2 l_3}^{PLS} = - \left\{ b_{l_3} c_{l_1} \frac{l_2(l_2+1) - l_1(l_1+1) - l_3(l_3+1)}{2} + cyc.perm. \right\} \sqrt{\frac{(2l_1+1)(2l_2+1)(2l_3+1)}{4\pi}} \begin{pmatrix} l_1 & l_2 & l_3 \\ 0 & 0 & 0 \end{pmatrix} \equiv b_{l_1 l_2 l_3} I_{l_1 l_2 l_3}, \quad (32)$$

where we have defined the reduced bispectra $b_{l_1 l_2 l_3}$, which is useful in certain context (Bartolo, Matarrese & Riotto 2006), and the additional geometrical factor, which originates from the integral involving three spherical harmonics over the entire sky:

$$I_{l_1 l_2 l_3} \equiv \sqrt{\frac{(2l_1+1)(2l_2+1)(2l_3+1)}{4\pi}} \begin{pmatrix} l_1 & l_2 & l_3 \\ 0 & 0 & 0 \end{pmatrix}. \quad (33)$$

The cross-correlation power-spectra appearing in the above expression denotes the coupling of lensing with a specific form of secondary non-anisotropy (see e.g. Cooray & Hu (2000)).

$$\langle \Theta_{l' m'} b_{lm} \rangle = b_l \delta_{ll'} \delta_{mm'}. \quad (34)$$

The bispectrum contains all the information at the three-point correlation function level and can be reduced to one-point skewness or the two-point collapsed correlation function or the associated power-spectra, the skew spectrum. Here we have considered the secondaries of the CMB, however the analysis holds if external tracers such as the radio galaxy surveys such as NVSS or 21cm observations are used instead. In the next few subsections we will discuss the problem of estimation of the skew spectrum in a nearly optimal way. This will lead to a discussion of development of optimal techniques in subsequent sections. We will also tackle the problem of joint estimation of several bispectra and associated estimation errors. As is known and will be discussed in the following sections that the estimation of CMB bispectra is similar and related to the case of lensing reconstruction of the CMB sky (Smith, Zahn & Dore 2007).

5 ESTIMATION OF SKEW SPECTRA

The problem of estimation of the skew spectrum is very similar to that of the primary CMB bispectrum. There has been a recent surge in activity in this area, driven by the claim of detection of non-gaussianity in the WMAP data release (see e.g. Yadav & Wandelt (2008); Yadav et al. (2008); Yadav, Komatsu & Wandelt (2007)). Different techniques were developed which introduce various weighting schemes in the harmonic domain to make the method optimal (i.e. saturates the Cramer-Rao bound). Maps are constructed by weighting the observed CMB sky with l -dependent weights obtained from inflationary theoretical models. These weighted maps are then used to compute one-point quantities which are generalisation of skewness and can be termed mixed skewness. These mixed skewness measures are useful estimators of f_{NL} parameters. A more general treatment was provided

in Smith & Zaldarriaga (2006); Smith, Zahn & Dore (2007); Smith, Senatore & Zaldarriaga (2009) who took into account mode-mode coupling in an exact way with the use of proper inverse-covariance weighting of harmonic modes.

Recent work by Munshi & Heavens (2009) has improved the situation by focussing directly on the skew spectrum. Their technique does not compress all the available information in the bispectrum into a single number but provides a power-spectrum which depends on the harmonic wavenumber l . This method has the advantage of being able to separate various contributions as they will have different dependence on l s, thus allowing an assessment of whether any non-gaussianity is primordial or not. In this section we compute the contaminating secondary bispectrum contributions from lensing-secondary coupling.

5.1 Bispectra without line of sight integration involving ISW-lensing RS-lensing and SZ-lensing

The study of the bispectrum related to secondary anisotropy (see Cooray & Seth (2000) for more details and analytical modelling based on halo model which we use here) is arguably as important as that generated by the primary anisotropy. Primary non-Gaussianity in simpler inflationary models is vanishingly small (Salopek & Bond 1990, 1991; Falk et al. 1993; Gangui et al. 1994; Acquaviva et al. 2003; Maldacena 2003); see Bartolo, Matarrese & Riotto (2006) and references therein for more details. However, variants of simple inflationary models such as multiple scalar fields (Linde & Mukhanov 1997; Lyth, Ungarelli & Wands 2003), features in the inflationary potential, non-adiabatic fluctuations, non-standard kinetic terms, warm inflation (Gupta, Berera & Heavens 2002; Moss & Xiong 2007), or deviations from Bunch-Davies vacuum can all lead to a much higher level of non-Gaussianity.

Early observational work on the bispectrum from COBE (Komatsu et al. 2002) and MAXIMA (Santos et al. 2003) was followed by much more accurate analysis with WMAP (Komatsu et al. 2003; Creminelli et al. 2007; Spergel et al. 2007). The primary bispectrum encodes information about inflationary dynamics and hence can constrain various inflationary scenarios, where as the secondary bispectrum will provide valuable information regarding the low-redshift universe and constrain structure formation scenarios. These bispectra are generated because of the cross-correlation effect of lensing due to various intervening materials and the secondary anisotropy such as the Sunyaev-Zeldovich effect due to inverse Compton scattering of CMB photons from hot gas in intervening clusters.

$$b_{l_1 l_2 l_3} = -\frac{1}{2} \left[\{ (l_2(l_2 + 1) - l_1(l_1 + 1) - l_3(l_3 + 1)) \} C_{l_1}^S b_{l_3} + \text{cyc.perm.} \right]. \quad (35)$$

The power spectrum C_l^S is the unlensed power spectrum of the CMB anisotropy. We have introduced the subscript S to distinguish it from the C_l s that appear in the denominator which take contribution from the instrumental noise from signal to noise computation point of view ($C_l = C_l + N/b(l)^2$, where N is the instrumental noise and $b(l)$ is the beam function in multipole space). We define the different fields which are constructed from underlying harmonics and corresponding C_l s. These will be useful for constructing an unbiased near optimal estimator.

$$\begin{aligned} A_{lm}^{(1)} &= \frac{a_{lm}}{C_l} C_l^S; & B_{lm}^{(1)} &= l(l+1) \frac{a_{lm}}{C_l}; & C_{lm}^{(1)} &= \frac{a_{lm}}{C_l} b_l \\ A_{lm}^{(2)} &= -l(l+1) \frac{a_{lm}}{C_l} C_l^S; & B_{lm}^{(2)} &= \frac{a_{lm}}{C_l}; & C_{lm}^{(2)} &= \frac{a_{lm}}{C_l} b_l \\ A_{lm}^{(3)} &= \frac{a_{lm}}{C_l} C_l^S; & B_{lm}^{(3)} &= \frac{a_{lm}}{C_l}; & C_{lm}^{(3)} &= l(l+1) \frac{a_{lm}}{C_l} b_l \end{aligned} \quad (36)$$

The corresponding fields that we construct are $A^{(i)}(\hat{\Omega}) \equiv \sum_{lm} Y_{lm}(\hat{\Omega}) A_{lm}^{(i)}$, and in an analogous manner $B^{(i)}$ and $C^{(i)}$. The optimised skew spectrum in the presence of all-sky coverage and homogeneous noise can now be written as:

$$C_l^{2,1} = \frac{1}{2l+1} \sum_m \sum_i \text{Real}\{ (A^{(i)}(\hat{\Omega}) B^{(i)}(\hat{\Omega}))_{lm} C^{(i)*}(\hat{\Omega})_{lm} \} + \text{cyc.perm.} \quad (37)$$

The cyclic terms that are considered here will have to be constructed likewise from the corresponding terms in the expression for the reduced bispectrum discussed above 35. The linear-order correction terms which needs to be included in the absence of spherical symmetry due to presence of cuts to avoid the galactic foreground and the inhomogeneous noise can be written as:

$$\hat{C}_l^{2,1} = \frac{1}{f_{sky}} \sum_i \left[\tilde{C}_l^{AB,C} - C_l^{A(B,C)} - C_l^{B(A,C)} - C_l^{(AB),C} \right]^{(i)} + \text{cyc.perm.} \quad (38)$$

The terms without averaging such as $\tilde{C}_l^{AB,C}$ are direct estimates from the observed partial sky with inhomogeneous noise. The Monte Carlo corrections such as $C_l^{A(B,C)}$ are constructed by cross-correlating the product of the observed map A and a Monte Carlo map B with a Monte Carlo map C and then taking an ensemble average over many realisations. The denominator f_{sky} , which represent the fraction of the sky covered, is introduced to correct for the effect of partial sky coverage. This is an approximate way to treat the mode-mode coupling due to partial sky coverage and known to be a good approximation for higher l . The skewness associated with this form of bispectra can be expressed as a weighted sum of the corresponding C_l s:

$$\hat{S} = \sum_l (2l+1) \hat{C}_l^{2,1} = \sum_{l_1 l_2} \frac{\hat{B}_{l_1 l_2} B_{l_1 l_2}}{C_l C_{l_1} C_{l_2}}. \quad (39)$$

Constructing such weighted maps clearly can be seen as a way to construct a matched-filter estimator for the detection of non-Gaussianity. It is optimally weighted by the inverse cosmic variance and achieves maximum response when the observed non-Gaussianity matches with a specific theoretical input. The skew spectrum also allows for analysis of more than one specific type of non-gaussianity from the same data - allowing a joint analysis to determine cross-contamination from various contributions.

5.2 Bispectra involving line of sight integration: The Ostriker-Vishniac effect and its correlation with other secondary anisotropies

Another set of secondary bispectra involving any of the Ostriker-Vishniac (e.g. see Jaffe & Kamionkowski (2004b)) effect, SZ thermal effect, or the kinetic SZ effect (Cooray 2001b) or a combination of these have the following form of reduced bispectrum (Cooray & Hu 2000), which involves a line of sight integration along r :

$$b_{l_1 l_2 l_3} = \int dr f_{l_1}(r) g_{l_2}(r) + \text{cyc.perm.} \quad (40)$$

The construction of weighted maps follow the same principle with the use of kernels $f_l(r)$ and $g_l(r)$ that are associated with any of the scattering secondaries that involve a line of sight integration. Numerical implementation of line of sight will naturally have to deal with an optimal method to include the quadrature. Defining

$$A_{lm}(r) = \frac{a_{lm}}{C_l} f_l(r); \quad B_{lm}(r) = \frac{a_{lm}}{C_l} g_l(r); \quad C_{lm}(r) = \frac{a_{lm}}{C_l}. \quad (41)$$

A and B are fields constructed from the generic function represented by f_l and g_l . Following Munshi & Heavens (2009) we construct

$$\hat{C}_l^{2,1}(r) = \frac{1}{2l+1} \int dr r \sum_m \text{Real}\{(AB)_{lm}(\hat{\Omega}, r) C(\hat{\Omega}, r)_{lm}\} + \text{cyc.perm.} \quad (42)$$

and from this compute the skew-spectrum:

$$\hat{C}_l^{2,1} = \int dr \hat{C}_l^{2,1}(r). \quad (43)$$

This is the generalisation of the all-sky estimator of the skew spectrum of Munshi & Heavens (2009), but for three distinct fields.

The corresponding one-point skewness can be written as

$$\hat{S} = \sum_l (2l+1) \hat{C}_l^{2,1} = \sum_{l_1 l_2} \frac{B_{l_1 l_2} \hat{B}_{l_1 l_2}}{C_l C_{l_2} C_{l_3}}. \quad (44)$$

In the next section we consider the contamination of the primary skew spectrum by secondary non-Gaussianity from point sources. As before we have absorbed the beam in the harmonic coefficients of the data vector a_{lm} . As before, C_l s also take contribution from the noise C_l^N as well as from the theoretical CMB powerspectra C_l^S , i.e. $C_l = C_l^S + C_l^N$

5.3 General Expression

From the examples above, its clear that from a very general consideration if the reduced bispectrum can be decomposed in such a way it consists of terms, which can be used to construct fields such as $A^{(i)}$, $B^{(i)}$ and $C^{(i)}$ (not necessarily of a specific form) a skew-spectrum can always be constructed by similar manipulation. In certain cases the $\hat{C}_l^{2,1}(r)$ might actually also have radial dependence, in which case a line of sight integration needs to be performed to match observations.

$$\hat{C}_l^{2,1} = \frac{1}{(2L+1)} \sum_{l' l''} \sum_{ij} \frac{(2l'+1)(2l''+1)}{144\pi} \begin{pmatrix} l' & l'' & L \\ 0 & 0 & 0 \end{pmatrix}^2 \frac{1}{C_L} \frac{1}{C_{l'}} \frac{1}{C_{l''}} [A_{l_1}^i B_{l_2}^i C_{l_3}^i + \text{cyc.perm.}] [A_{l_1}^j B_{l_2}^j C_{l_3}^j + \text{cyc.perm.}] \quad (45)$$

5.4 Cross-contamination from Point Sources and Primary non-Gaussianity

The bispectra associated with point sources is modelled as $b_{l_1 l_2 l_3}^{ps} = \text{const.}$ The constant depends on the flux limit. More complicated modelling which incorporates certain aspects of halo models can be used for better accuracy Serra & Cooray (2008).

$$S = \sum_{l_1 l_2 l_3} \frac{B_{l_1 l_2 l_3}^{ps} B_{l_1 l_2 l_3}^{sec}}{C_{l_1} C_{l_2} C_{l_3}}. \quad (46)$$

Similarly given a model of primary non-Gaussianity one can construct a theoretical model for computation of $B_{l_1 l_2 l_3}^{prim}$ (see Munshi & Heavens 2009 for more about various models and construction of optimal estimators). While study of primary non-Gaussianity is important in its own right for the study of secondaries they can confuse the study.

$$S = \sum_{l_1 l_2 l_3} \frac{B_{l_1 l_2 l_3}^{prim} B_{l_1 l_2 l_3}^{sec}}{C_{l_1} C_{l_2} C_{l_3}} \quad (47)$$

Similar results hold at the level of the skew spectrum. A more general treatment based on Fisher analysis of multiple bispectra is presented in the subsequent sections.

In addition to various sources mentioned above, second-order corrections to the gravitational potential through gravitational instability too can also act as a source of secondary non-Gaussianity (Munshi, Souradeep & Starobinsky 1995).

6 OPTIMISED ANALYSIS OF MIXED BISPECTRA

Starting from Babich (2005) a complete analysis of bispectrum in the presence of partial sky coverage and inhomogeneous noise was developed by various authors (Babich 2005; Creminelli et al. 2006; Yadav et al. 2008). A specific form for a bispectrum estimator was introduced which is both unbiased and optimal. This was further developed and used by Smith, Zahn & Dore (2007) for lensing reconstruction and by Smith & Zaldarriaga (2006) for general bispectrum analysis. for one-point estimator for f_{NL} . The analysis depends on finding suitable inverse cosmic variance weighting C^{-1} of modes. It deals with mode-mode coupling in an exact way. In a recent work Munshi & Heavens (2009) further extended this analysis by incorporating two-point statistics or the skew spectrum which we have already introduced above. We generalise their results in this work for the case of mixed bispectra for the case of both one-point and two-point studies involving three-way correlations. The analytical results presented here are being kept as general as possible. However in the next sections we specialise them to individual cases of lensing reconstruction and the mixed bispectrum associated with lensing and the SZ effect as concrete examples.

6.1 One-point Estimator: Mixed Skewness $\langle X(\hat{\Omega})Y(\hat{\Omega})Z(\hat{\Omega}) \rangle$

We are interested in constructing an optimal and unbiased estimator for the estimation of mixed skewness $\langle X(\hat{\Omega})Y(\hat{\Omega})Z(\hat{\Omega}) \rangle$. The fields $X(\Omega) = \sum_{lm} X_{lm} Y_{lm}(\hat{\Omega})$ and similarly for Y and Z , are defined over the entire sky, though observed with a mask and nonuniform noise coverage. The non-uniform coverage imprints a mode-mode coupling $[C_{XX}]_{l_1 m_1, l_2 m_2}^{-1}$ in the observed multipoles of a given field in the harmonic space $\langle X_{l_1 m_1} X_{l_2 m_2} \rangle$. For the construction of the optimal estimator it will be useful to define $\tilde{X}_{l_1 m_1}$ as

$$\tilde{X}_{l_1 m_1} \equiv [C_{XX}]_{l_1 m_1, l_a m_a}^{-1} X_{l_a m_a}. \quad (48)$$

Here $\tilde{X}_{lm}$ represents the harmonics of the data X with inverse covariance weighting. Next we need to deal with the covariance matrix of the modes $\tilde{X}_{l_1 m_1}$ in terms of that of $X_{l_1 m_1}$. The auto covariance matrix for X , C_{XX} , and that of $\tilde{X}$, $\tilde{C}_{XX}$ are related by the following expression:

$$[\tilde{C}_{XX}]_{l_a m_a, l_b m_b} \equiv \langle \tilde{X}_{l_a m_a} \tilde{X}_{l_b m_b} \rangle = [C_{XX}^{-1}]_{l_a m_a, l_b m_b}. \quad (49)$$

Similarly, the cross-covariance for two different fields $\tilde{X}$ and $\tilde{Y}$ with inverse variance weighting, in harmonic space can be written as:

$$\tilde{C}_{XY} \equiv \langle \tilde{X}_{l_a m_a} \tilde{Y}_{l_b m_b} \rangle = C_{XX}^{-1} C_{XY} C_{YY}^{-1}. \quad (50)$$

The estimator that we construct will be based on functions $\hat{Q}[\tilde{X}, \tilde{Y}, \tilde{Z}]$ which depends on the input fields, and its derivatives w.r.t. the fields e.g. $\partial \hat{Q}[\tilde{X}, \tilde{Y}] / \partial \tilde{Z}_{lm}$. The derivatives are themselves a map with harmonics described by the free indices lm , and are constructed out of two other maps. The function $\hat{Q}$ on the other hand is an ordinary number which depends on all three input functions and lacks free indices.

$$\hat{Q}[\tilde{X}, \tilde{Y}, \tilde{Z}] \equiv \frac{1}{6} \sum_{lm'l'm''} B_{ll'l''}^{XYZ} \begin{pmatrix} l & l' & l'' \\ m & m' & m'' \end{pmatrix} \tilde{X}_{lm} \tilde{Y}_{l'm'} \tilde{Z}_{l''m''} \quad (51)$$

$$\frac{\partial \hat{Q}[\tilde{X}, \tilde{Y}]}{\partial \tilde{Z}_{lm}} \equiv \sum_{lm'l'm''} B_{ll'l''}^{XYZ} \begin{pmatrix} l & l' & l'' \\ m & m' & m'' \end{pmatrix} \tilde{X}_{l'm'} \tilde{Y}_{l''m''}. \quad (52)$$

Similar expressions hold for other fields such as $\tilde{X}_{lm}$, $\tilde{Y}_{lm}$. Introducing a more compact notation x_i , where $x_1 = X$, $x_2 = Y$, $x_3 = Z$ we can write the one-point estimator for the mixed skewness as:

$$\hat{E}[\tilde{x}_i] = \frac{1}{F} \left\{ Q[\tilde{x}_i] - \sum_i [\tilde{x}_i]_{l_a m_a}^i \langle \partial_{l_a m_a}^i Q[\tilde{x}_i] \rangle \right\}. \quad (53)$$

This is a main result of the paper, generalising work by Smith & Zaldarriaga (2006) to mixed fields.

The ensemble averaging $\langle \rangle$ in the linear terms represents Monte-Carlo averaging using simulated non-Gaussian maps. The associated Fisher matrix (a scalar in this case) can be written in terms of the functions $Q[\tilde{x}_i]$, its derivative and the cross-covariance matrices involving different fields.

$$F = \langle \partial_{l_a m_a}^i Q[\tilde{x}_i] [C^{-1}]_{l_a m_a, l_b m_b}^{ij} \partial_{l_b m_b}^j Q[\tilde{x}_j] \rangle - \langle \partial_{l_a m_a}^i Q[\tilde{x}_i] [C^{-1}]_{l_a m_a, l_b m_b}^{ij} \langle \partial_{l_b m_b}^j Q[\tilde{x}_j] \rangle \rangle. \quad (54)$$

Here we have used the shorthand notation for $\langle x_{l_a m_a}^i x_{l_b m_b}^j \rangle = [C^{-1}]_{l_a m_a, l_b m_b}^{ij}$. In case of joint estimation of different bispectra from the same data sets we can extend the above discussion and write:

$$E[\tilde{x}_i] = F^{-1}_{\alpha\beta} \left\{ Q^\beta[\tilde{x}_i] - [\tilde{x}_i]_{l_a m_a}^i \langle \partial_{l_a m_a}^i Q^\beta[\tilde{x}_i] \rangle \right\}. \quad (55)$$

Here the Fisher matrix $F_{\alpha\beta}$ encodes the inverse estimator covariance for different mixed bispectra B^α and B^β , α and β represents different types of bispectra recovered using the same data sets.

$$F_{\alpha\beta} = \sum B_{l_1 l_2 l_3}^\alpha B_{l_4 l_5 l_6}^\beta \left[[\tilde{C}^{XX}]_{l_1 m_1, l_4 m_4} [\tilde{C}^{YY}]_{l_2 m_2, l_5 m_5} [\tilde{C}^{ZZ}]_{l_3 m_3, l_6 m_6} + [\tilde{C}^{XX}]_{l_1 m_1, l_4 m_4} [\tilde{C}^{YZ}]_{l_2 m_2, l_6 m_6} [\tilde{C}^{ZY}]_{l_3 m_3, l_5 m_5} + [\tilde{C}^{XY}]_{l_1 m_1, l_5 m_5} [\tilde{C}^{YZ}]_{l_2 m_2, l_6 m_6} [\tilde{C}^{ZX}]_{l_3 m_3, l_4 m_4} + [\tilde{C}^{XZ}]_{l_1 m_1, l_6 m_6} [\tilde{C}^{ZY}]_{l_3 m_3, l_5 m_5} [\tilde{C}^{YX}]_{l_2 m_2, l_4 m_4} \right]. \quad (56)$$




Figure 4. The cross-spectra b_l introduced in Eq.(32) required for the construction of the bispectra is plotted for ISW cross lensing (right panel) and SZ cross lensing (left panel) bispectrum as a function of l . See text for details.

The cyclic permutations here represent two additional terms with permutations of super-scripts X, Y, Z along with associated subscripts. The Fisher matrix (which is a number in this particular case) for the mixed bispectrum in case of all-sky coverage and constant variance noise can be expressed as:

$$F = \frac{1}{6} \sum_{l_1 l_2 l_3} B_{l_1 l_2 l_3}^{XYZ} B_{l_1 l_2 l_3}^{XYZ} \left[\frac{1}{C_{l_1}^{XX}} \frac{1}{C_{l_2}^{YY}} \frac{1}{C_{l_3}^{ZZ}} + \frac{1}{C_l^{XX}} \left(\frac{C_{l_1}^{YZ}}{C_{l_2}^{XX} C_{l_3}^{YY}} \right)^2 + 2 \frac{C_{l_1}^{YZ}}{C_{l_2}^{ZZ} C_{l_3}^{YY}} \frac{C_{l_1}^{ZX}}{C_{l_2}^{XX} C_{l_3}^{XX}} \frac{C_{l_1}^{XY}}{C_{l_2}^{XX} C_{l_3}^{YY}} \right] \quad (57)$$

6.1.1 Special Case (A): $Z=Y, \langle X(\hat{\Omega})Y^2(\hat{\Omega}) \rangle$

In certain practical situations we will encounter cases where two of the three fields are identical. The corresponding Fisher matrix can be recovered by simply setting $Z = Y$.

$$F = \frac{1}{6} \sum_{l_1 l_2 l_3} \left[2 B_{l_1 l_2 l_3}^{XYY} B_{l_1 l_2 l_3}^{XYY} \left(\frac{1}{C_{l_1}^{XX}} \right) \left(\frac{1}{C_{l_2}^{YY}} \right)^2 + 2 B_{l_1 l_2 l_3}^{XYY} B_{l_3 l_2 l_1}^{XYY} \left(\frac{C_{l_1}^{XY}}{C_{l_1}^{XX} C_{l_1}^{YY}} \right)^2 \left(\frac{1}{C_{l_2}^{YY}} \right) \right]. \quad (58)$$

6.1.2 Special Case (B): $Z=Y=X, \langle X^3(\hat{\Omega}) \rangle$

Finally, if we identify all three fields to recover the case ordinary or pure bispectrum corresponding to the case $X = Y = Z$.

$$F_{\alpha\beta} = \frac{1}{6} \sum B_{l_1 l_2 l_3}^{\alpha} B_{l_4 l_5 l_6}^{\beta} [C^{-1}]_{l_1 m_1, l_4 m_4} [C^{-1}]_{l_2 m_2, l_5 m_5} [C^{-1}]_{l_3 m_3, l_6 m_6}. \quad (59)$$

In the limit of all-sky coverage and constant variance noise the estimator reduces to:

$$F_{\alpha\beta} = \frac{1}{6} \sum_{l_1 l_2 l_3} \frac{B_{l_1 l_2 l_3}^{XYZ} B_{l_1 l_2 l_3}^{XYZ}}{C_{l_1} C_{l_2} C_{l_3}} \quad (60)$$

For high ls a scaling $f_{sky}^{-1/2}$ is sufficient to describe the effect of partial sky coverage on the error covariance matrix.

6.2 Two-point Estimators: Mixed skew spectrum

We begin by constructing the functions Q_L and its derivative w.r.t. various input fields. We use these to construct an optimal and unbiased estimator to correlate the field $X(\hat{\Omega})$ with the product of two such fields $Y(\hat{\Omega})Z(\hat{\Omega})$. We consider the most general possible case of the skew spectrum associated with the mixed bispectrum B^{XYZ} .

$$\hat{Q}_L[\tilde{X}, \tilde{Y}, \tilde{Z}] \equiv \sum_M \tilde{X}_{LM} \sum_{l'm', l''m''} B_{Ll'l''} \begin{pmatrix} L & l' & l'' \\ M & m' & m'' \end{pmatrix} \tilde{Y}_{l'm'} \tilde{Z}_{l''m''} \quad (61)$$

$$\partial_{lm}^X \hat{Q}_L[\tilde{Y}, \tilde{Z}] \equiv \delta_{Ll} \sum_{l'm', l''m''} B_{Ll'l''} \begin{pmatrix} L & l' & l'' \\ m & m' & m'' \end{pmatrix} \tilde{Y}_{l'm'} \tilde{Z}_{l''m''} \quad (62)$$

$$\partial_{lm}^Y \hat{Q}_L[\tilde{X}, \tilde{Z}] \equiv \sum_M \tilde{X}_{LM} \sum_{l'm'} B_{Ll'l'} \begin{pmatrix} L & l & l' \\ M & m & m' \end{pmatrix} \tilde{Z}_{l'm'}; \quad \partial_{lm}^Z \hat{Q}_L[\tilde{X}, \tilde{Y}] \equiv \sum_M \tilde{X}_{LM} \sum_{l'm'} B_{Ll'l'} \begin{pmatrix} L & l & l' \\ M & m & m' \end{pmatrix} \tilde{Y}_{l'm'}. \quad (63)$$

12 Munshi, Valageas, Cooray & Heavens

While Q_L is a number (cubic function of input maps) for a given L , the derivatives are maps which are quadratic in the input maps. The derivatives will be important in constructing the linear terms which are important in reducing the variance of the estimator in an absence of spherical symmetry, which is the case in the presence of inhomogeneous noise or partial sky coverage. Using these expressions we can write down the optimised bispectra

$$\hat{E}_L^{X,YZ}[x_i] = [N^{-1}]_{LL'}^{X,YZ} \left\{ Q_{L'}[\tilde{x}_i] - \sum_{i=1,2,3} [\tilde{x}_i]_{lm} \langle \partial_{lm}^i Q_{L'}[\tilde{x}_i] \rangle_{MC} \right\}. \quad (64)$$

The normalisation matrix $N_{LL'}$ is related to the Fisher matrix $F_{LL'} = N_{LL'}^{-1}$, and can be expressed as:

$$N_{LL'}^{X,YZ} = \frac{1}{3} \langle \{ \partial_{l_1 m_1} Q_L[\tilde{x}] \} [C_{l_1 m_1, l_2 m_2}^{-1}]^{ij} \{ \partial_{l_2 m_2}^j Q_{L'}[\tilde{x}] \} \rangle - \frac{1}{3} \{ \langle \partial_{l_1 m_1}^i Q_L[\tilde{x}] \rangle [C_{l_1 m_1, l_2 m_2}^{-1}]^{ij} \{ \langle \partial_{l_2 m_2}^j Q_{L'}[\tilde{x}] \rangle \}. \quad (65)$$

Finally the Fisher matrix can be written as:

$$\begin{aligned} F_{LL'} = & \sum_{MM'} \sum_{l_i l'_i m_i m'_i} B_{L l_1 l'_1}^{XYZ} B_{L' l'_2 l_2}^{XYZ} \begin{pmatrix} L & l_1 & l'_1 \\ M & m_1 & m'_1 \end{pmatrix} \begin{pmatrix} L' & l_2 & l'_2 \\ M' & m_2 & m'_2 \end{pmatrix} \\ & \times \frac{1}{6} \{ [\tilde{C}^{XX}]_{LM, L'M'} [\tilde{C}^{YY}]_{l_1 m_1, l'_1 m'_1} [\tilde{C}^{ZZ}]_{l_2 m_2, l'_2 m'_2} + [\tilde{C}^{XY}]_{LM, l'_1 m'_1} [\tilde{C}^{YZ}]_{l_1 m_1, L'M'} [\tilde{C}^{ZX}]_{l_2 m_2, l'_2 m'_2} + \\ & [\tilde{C}^{XZ}]_{LM, l'_2 m'_2} [\tilde{C}^{ZY}]_{l_2 m_2, l'_1 m'_1} [\tilde{C}^{YX}]_{l_1 m_1, L'M'} + [\tilde{C}^{XX}]_{LM, L'M'} [\tilde{C}^{ZZ}]_{l_2 m_2, l'_1 m'_1} [\tilde{C}^{ZY}]_{LM, l'_2 m'_2} + \\ & [\tilde{C}^{YY}]_{l_1 m_1, l'_1 m'_1} [\tilde{C}^{XZ}]_{LM, l'_2 m'_2} [\tilde{C}^{ZX}]_{l_2 m_2, L'M'} + [\tilde{C}^{ZZ}]_{l_2 m_2, l'_2 m'_2} [\tilde{C}^{XY}]_{LM, l'_1 m'_1} [\tilde{C}^{YX}]_{l_1 m_1, L'M'} \}. \end{aligned} \quad (66)$$

In the case of near all-sky experiments the off-diagonal elements of the Fisher matrix will be relatively smaller. The diagonal elements as before can be scaled by f_{sky} (the fraction of the sky covered by a near all-sky experiment). The covariance matrices can now be expressed only as a function of the related C_i s the auto- and cross-correlation power spectra:

$$\begin{aligned} F_{LL'} = & \delta_{LL'} \sum_{ll'} \left\{ B_{L ll'}^{XYZ} B_{L l l'}^{XYZ} \frac{1}{C_L^{XX}} \frac{1}{C_l^{YY}} \frac{1}{C_{l'}^{ZZ}} + B_{L ll'}^{XYZ} B_{L l l'}^{XYZ} \frac{1}{C_L^{XX}} \left(\frac{C_l^{YZ}}{C_l^{YY} C_l^{ZZ}} \right) \left(\frac{C_{l'}^{YZ}}{C_{l'}^{YY} C_{l'}^{ZZ}} \right) \right\} \\ & + \sum_l \left\{ B_{L' l L}^{XYZ} B_{L l L'}^{XYZ} \left(\frac{1}{C_l^{YY}} \right) \left(\frac{C_{L'}^{XZ}}{C_L^{XX} C_{L'}^{ZZ}} \right) \left(\frac{C_L^{ZX}}{C_L^{XX} C_L^{ZZ}} \right) + B_{L L' l}^{XYZ} B_{L' l L}^{XYZ} \left(\frac{1}{C_{L'}^{ZZ}} \right) \left(\frac{C_L^{XY}}{C_L^{XX} C_L^{YY}} \right) \left(\frac{C_{L'}^{XY}}{C_{L'}^{XX} C_{L'}^{YY}} \right) \right. \\ & \left. + B_{L L' l}^{XYZ} B_{L l L'}^{XYZ} \left(\frac{C_L^{XY}}{C_L^{XX} C_L^{YY}} \right) \left(\frac{C_{L'}^{YZ}}{C_{L'}^{YY} C_{L'}^{ZZ}} \right) \left(\frac{C_L^{ZX}}{C_L^{XX} C_L^{ZZ}} \right) + B_{L l L'}^{XYZ} B_{L' l L}^{XYZ} \left(\frac{C_L^{XZ}}{C_L^{XX} C_L^{ZZ}} \right) \left(\frac{C_{l'}^{ZY}}{C_{l'}^{YY} C_{l'}^{ZZ}} \right) \left(\frac{C_{L'}^{YX}}{C_{L'}^{XX} C_{L'}^{YY}} \right) \right\}. \end{aligned} \quad (67)$$

The other two terms represented by cyc.perm. consist of terms with suitable permutations of superscripts X, Y and Z . In deriving these expressions all-sky limits of $\tilde{C}_{lm, l' m'}^{XX} = (1/C_l^{XX}) \delta_{ll'} \delta_{mm'}$ were used along with the fact that we can write $\tilde{C}_{lm, l' m'}^{XY} = \{C_l^{XY}/C_l^{XX} C_l^{YY}\} \delta_{ll'} \delta_{mm'}$ for all sky case. For the case when the cross-correlation among two or more fields vanish the expression simplifies considerably.

In case of joint estimation of several bispectra from the same data one can write the following expression:

$$\hat{E}_L^\alpha[x_i] = \sum_{L'} \sum_{\beta} [N^{-1}]_{LL'}^{\alpha\beta} \left\{ Q_{L'}^\beta[\tilde{x}_i] - \sum_{i=1,2,3} [\tilde{x}_i]_{lm} \langle \partial_{lm} Q_{L'}^\beta[\tilde{x}_i] \rangle_{MC} \right\}. \quad (68)$$

Here the indices α and β correspond to various power spectra $[B_{l_1 l_2 l_3}^{XYZ}]^\alpha$ or $[B_{l_1 l_2 l_3}^{XYZ}]^\beta$ which are associated with bispectra that can be jointly estimated from the same data $[\tilde{x}_i]$. Below we consider two special cases for the skew spectra that we have considered so far. The expressions for $F_{LL'}^{\alpha\beta}$ and $N_{LL'}^{\alpha\beta}$ can be obtained simply by replacing the product $B_{L l_1 l'_1}^{XYZ} B_{L l'_1 l_2}^{XYZ}$ by $[B_{l_1 l_2 l_3}^{XYZ}]^\alpha [B_{l_1 l_2 l_3}^{XYZ}]^\beta$. In certain situation when accurate noise modelling is difficult or unlikely an approximate proxy for C^{-1} is used in the form of a regularization matrix R which acts as a smoothing of the data. The resulting data vector $\tilde{x}_i^R = R x_i$ is now used for developing an unbiased but suboptimal estimator by replacing $\tilde{x}_i$ with $\tilde{x}_i^R$.

6.2.1 Special Case (A): $Z=Y$

The estimator in this case corresponds to E_l^{X,Y^2} .

$$\begin{aligned} F_{LL'} = & 2\delta_{LL'} \sum_{ll'} \left\{ B_{L ll'}^{XYY} B_{L l l'}^{XYY} \frac{1}{C_L^{XX}} \frac{1}{C_l^{YY}} \frac{1}{C_{l'}^{YY}} \right\} + \sum_l \left\{ 2 B_{L L' l}^{XYY} B_{L' l L}^{XYY} \left(\frac{C_L^{XY}}{C_L^{XX} C_L^{YY}} \right) \left(\frac{C_{L'}^{XY}}{C_{L'}^{XX} C_{L'}^{YY}} \right) \frac{1}{C_l^{YY}} \right. \\ & \left. + B_{L L' l}^{XYY} B_{L l L'}^{XYY} \frac{1}{C_{L'}^{YY}} \left(\frac{C_L^{XY}}{C_L^{XX} C_L^{YY}} \right) \left(\frac{C_{l'}^{YX}}{C_{l'}^{XX} C_{l'}^{YY}} \right) + B_{L l L'}^{XYY} B_{L' l L}^{XYY} \frac{1}{C_l^{YY}} \left(\frac{C_L^{XY}}{C_L^{XX} C_L^{YY}} \right) \left(\frac{C_{L'}^{YX}}{C_{L'}^{XX} C_{L'}^{YY}} \right) \right\}. \end{aligned} \quad (69)$$




Figure 5. The cross-spectra b_l introduced in Eq.(32) required for the construction of the bispectra is plotted for point source(PS) cross lensing (right panel), ISW cross lensing (middle panel) and SZ cross lensing (left panel) bispectrum as a function of l . See text for details.

6.2.2 Special Case (B): $Z=Y=X$

$$F_{LL'} = 2\delta_{LL'} \sum_{ll'} \left\{ B_{LL'l}^{XXX} B_{LL'l'}^{XXX} \frac{1}{c_L^{XX}} \frac{1}{c_l^{XX}} \frac{1}{c_{l'}^{XX}} \right\} + \sum_l \left\{ B_{LL'l}^{XXX} B_{LL'l}^{XXX} \frac{1}{c_L^{XX}} \frac{1}{c_{l'}^{XX}} \frac{1}{c_l^{XX}} \right\} \quad (70)$$

6.2.3 Special Case (C): $Z=X$

The estimator we consider here is $E_l^{X,XY}$ can not derived by simple identification of superscript. We define the new functions related to the estimators Q_L , $\partial_{lm}^X \hat{Q}_L$ and $\partial_{lm}^Y \hat{Q}_L$ according to the same prescription above.

$$\hat{Q}_L[\tilde{X}, \tilde{Y}] \equiv \sum_M \tilde{X}_{LM} \sum_{l'm', l''m''} B_{Ll'l''} \begin{pmatrix} L & l' & l'' \\ M & m' & m'' \end{pmatrix} \tilde{X}_{l'm'} \tilde{Y}_{l''m''} \quad (71)$$

$$\partial_{lm}^Y \hat{Q}_L[\tilde{X}, \tilde{Y}] \equiv \sum_M \tilde{X}_{LM} \sum_{l'm', l''m''} B_{Ll'l''} \begin{pmatrix} L & l' & l'' \\ m & m' & m'' \end{pmatrix} \tilde{X}_{l'm'} \quad (72)$$

$$\partial_{lm}^X \hat{Q}_L[\tilde{X}, \tilde{Y}] \equiv \sum_{LM} \tilde{X}_{LM} \sum_{l'm'} B_{Ll'l'} \begin{pmatrix} L & l & l' \\ M & m & m' \end{pmatrix} \tilde{X}_{l'm'} + \delta_{Ll} \delta_{Mm} \sum_{l'm'} B_{Ll'l'} \begin{pmatrix} L & l & l' \\ M & m & m' \end{pmatrix} \tilde{X}_{l'm'} \tilde{Y}_{l'm'}. \quad (73)$$

The estimator in this case takes the form:

$$\hat{E}_L[x_i] = [N^{-1}]_{LL'} \left\{ Q_{L'}[\tilde{x}_i] - \sum_{i=1,2} [\tilde{x}_i]_{lm} \langle \partial_{lm}^i Q_{L'}[\tilde{x}_i] \rangle_{MC} \right\} \quad (74)$$

$$F_{LL'} = \delta_{LL'} \sum_{ll'} \left\{ B_{LL'l}^{XYX} B_{LL'l'}^{XYX} \frac{1}{c_L^{XX}} \frac{1}{c_l^{XX}} \frac{1}{c_{l'}^{XX}} \right\} + \left\{ B_{LL'l}^{XYX} B_{LL'l'}^{XYX} \frac{1}{c_L^{XX}} \left(\frac{c_l^{XY}}{c_l^{XX} c_l^{YY}} \right) \left(\frac{c_{l'}^{XY}}{c_{l'}^{XX} c_{l'}^{YY}} \right) \right\} \quad (75)$$

$$+ \sum_l \left\{ B_{LL'l}^{XYX} B_{LL'l}^{XYX} \left(\frac{c_L^{XY}}{c_L^{XX} c_L^{YY}} \right) \left(\frac{c_l^{XY}}{c_l^{XX} c_l^{YY}} \right) \frac{1}{c_l^{XX}} + B_{LL'l}^{XYX} B_{LL'l}^{XYX} \frac{c_L^{XY}}{c_L^{XX} c_L^{YY}} \frac{c_l^{XY}}{c_l^{XX} c_l^{YY}} \frac{1}{c_{l'}^{XX}} \right. \\ \left. + B_{LL'l}^{XYX} B_{LL'l}^{XYX} \frac{1}{c_L^{XX}} \left(\frac{c_{l'}^{XY}}{c_{l'}^{XX} c_{l'}^{YY}} \right) \left(\frac{c_l^{XY}}{c_l^{XX} c_l^{YY}} \right) + B_{LL'l}^{XYX} B_{LL'l}^{XYX} \frac{1}{c_L^{XX}} \frac{1}{c_{l'}^{XX}} \frac{1}{c_l^{YY}} \right\} \quad (76)$$

7 SPECIFIC EXAMPLES

The discussion so far has been completely general. We specialise now for a few practical cases of cosmological importance. These correspond to the study of mixed bispectra associated with lensing induced correlation of secondaries and CMB as well as frequency cleaned SZ catalogs against CMB sky.

7.1 Lensing Reconstruction

7.1.1 One-point estimator:

Various estimators associated with lensing reconstruction were introduced by different authors, e.g. (Hu 2000; Hu & Okamoto 2002)). It was recently studied by Smith, Zahn & Dore (2007) and was used to probe effect of lensing in CMB by cross-correlating with external data-set such as NVSS survey against WMAP observations.

$$S^{lens} = \frac{1}{2N} \sum_{l_i m_i} B_{l_1 l_2 l_3}^{\delta\delta\psi} [\tilde{\delta}_{l_1 m_1} \tilde{\delta}_{l_2 m_2} \tilde{\psi}_{l_3 m_3} - [C^{\phi\phi}]_{l_1 m_1, l_2 m_2}^{-1} \tilde{\psi}_{l_3 m_3}]. \quad (77)$$

This is achieved by writing the reconstructed lensing potential in terms of the CMB harmonics and cross-correlating it with low-redshift large-scale tracers such as galaxy surveys (Smith, Zahn & Dore 2007). The bispectrum $B_{l_1 l_2 l_3}^{\delta\delta\psi}$ depends in addition to the C_l s of the CMB multipole, on the cross-correlation between the CMB sky $\delta(\hat{\Omega})$ and the low-redshift tracer field $\psi(\hat{\Omega})$. The reduced bispectrum of interest $b_{l_1 l_2 l_3}^{\delta\delta\psi}$ and the related form factor $f_{l_1 l_2 l_3}$ can be written as:

$$b_{l_1 l_2 l_3}^{\delta\delta\psi} = \left\{ f_{l_1 l_2 l_3} C_{l_2}^{\delta\delta} + f_{l_2 l_1 l_3} C_{l_1}^{\delta\delta} \right\} C_{l_3}^{\delta\psi} \\ f_{l_1 l_2 l_3} = \frac{1}{2} \left\{ l_2(l_2 + 1) + l_3(l_3 + 1) - l_1(l_1 + 1) \right\} \quad (78)$$

The multipole $\delta_{l_1 m_1}$ and $\delta_{l_2 m_2}$ are associated with the CMB sky and $\psi_{l_3 m_3}$ is the multipole associated with the large-scale structure tracer at low redshift and hence correlates with the lensing potential (e.g. NVSS survey). The above estimator directly probes the cross-correlation between the lensing potential harmonics ϕ_{lm} constructed from temperature harmonics δ_{lm} and the harmonics of the tracers ψ_{lm} . It is interesting to notice that the estimator constructed lacks the term which signifies the correlation between $\delta(\hat{\Omega})$ and $\psi(\hat{\Omega})$ through the coupling $C^{\delta\psi}$. Though the bispectrum itself depends directly on the cross-power spectra. Using the results derived before we can write the Fisher matrix associated with this estimator can be written as:

$$F = N^{-1} = \frac{1}{2} \sum_{l_i m_i} B_{l_1 l_2 l_3}^{\delta\delta\psi} B_{l_4 l_5 l_6}^{\delta\delta\psi} [C^{\delta\delta}]_{l_1 m_1, l_4 m_4}^{-1} [C^{\delta\delta}]_{l_2 m_2, l_5 m_5}^{-1} [C^{\psi\psi}]_{l_3 m_3, l_6 m_6}^{-1}. \quad (79)$$

7.1.2 Estimators for the skew spectrum

If instead of the one-point estimator described above, we compute the two-point estimator or the skew spectrum as follows:

$$E_L[\tilde{\delta}, \tilde{\psi}] = [N^{-1}]_{LL'} \left\{ Q_{L'}[\tilde{\delta}, \tilde{\psi}] - \tilde{\psi} \langle \partial_{lm}^{\tilde{\psi}} Q_{L'}[\tilde{\delta}, \tilde{\psi}] \rangle \right\}. \quad (80)$$

The corresponding expressions for the functions $Q_{L'}[\tilde{\phi}]$ and $\partial_{lm} Q_{L'}[\tilde{\phi}, \tilde{\psi}]$ are given by:

$$Q_L = \sum_M \tilde{\psi}_{LM} \sum_{lm, l'm'} \tilde{\delta}_{lm} \tilde{\delta}_{l'm'} \begin{pmatrix} L & l & l' \\ m & m & m' \end{pmatrix}; \quad \partial_{lm}^{\psi} Q_L = \sum_{LM} \psi_{LM} \sum_{l'm'} B_{Ll'l'm'}^{\delta\delta\psi} \phi_{l'm'} \begin{pmatrix} L & l & l' \\ m & m & m' \end{pmatrix}. \quad (81)$$

Corresponding Fisher matrices can which turns out to be diagonal can be written as:

$$F_{LL'} = N_{LL'}^{-1} = \left\{ \langle \partial_{lm}^{\psi} Q_L [C^{\psi\psi}]_{lm, l'm'} \partial_{l'm'}^{\psi} Q_{L'} \rangle - \langle \partial_{lm}^{\psi} Q_L [C^{\psi\psi}]_{lm, l'm'} \rangle \langle \partial_{l'm'}^{\psi} Q_{L'} \rangle \right\}; \quad (82)$$

which finally leads us to:

$$F_{LL'} = \frac{1}{2} \sum_{ll'} B_{l_1 l_2 L}^{\delta\delta\psi} B_{l_3 l_4 L'}^{\delta\delta\psi} [C^{\phi\phi}]_{l_1 m_1, l_3 m_3}^{-1} [C^{\phi\phi}]_{l_2 m_2, l_4 m_4}^{-1} [C^{\psi\psi}]_{L M, L' M'}^{-1}. \quad (83)$$

In the limit of all sky survey and homogeneous noise we can write:

$$F_{LL'} = \delta_{LL'} \sum_{ll'} B_{ll'L}^{\delta\delta\psi} B_{ll'L'}^{\delta\delta\psi} \left(\frac{1}{C_L^{\phi\phi}} \right) \left(\frac{1}{C_l^{\phi\phi}} \right) \left(\frac{1}{C_{l'}^{\psi\psi}} \right) \quad (84)$$

A comparison with the previous estimator shows the presence of off-diagonal entries in the Fisher matrix even if direct correlation between δ and ψ are absent in the estimator.

It is possible to work with CMB sky without external data sets (such as NVSS or other galaxy surveys) to probe weak lensing e.g. the power spectrum of the lensing potential itself is related to four-point statistics of the temperature - which makes it noise dominated. use of external tracers such as galaxy surveys can reduce the problem to three-point level thus lowering the need on sensitivity of the instrument. The discussion above can have direct relevance for use of other tracers such as the one with neutral hydrogen observations (Zahn & Zaldarriaga 2006).




Figure 6. The cross-spectra b_l introduced in Eq.(32) required for the construction of the bispectra is plotted for ISW cross lensing (right panel) and SZ cross lensing (left panel) bispectrum as a function of l . See text for details.

7.2 Sunyaev-Zeldovich-CMB² mixed bispectrum

The secondary bispectrum caused by the Sunyaev-Zeldovich effect is one of the most pronounced secondary bispectrum among many others (Goldberg & Spergel 1999; Spergel & Goldberg 1999; Cooray & Hu 2000; Cooray 2000, 2001a,b). Following Cooray, Hu & Tegmark (2000) we study if frequency cleaned maps of all-sky CMB and SZ maps can also be used to construct power-spectra associated with the mixed bispectra with signal-to-noise ratio that can be detectable with ongoing CMB experiments. It probes mode-coupling effects generated by correlation involved with gravitational lensing angular deflections in CMB and the SZ effects due to large-scale pressure fluctuations. As before the estimator which can be constructed from the CMB $\tilde{a}_{lm}$ and the Sunyaev-Zeldovich s_{lm} multipoles. There is a possibility of constructing the correlating the product map $a(\hat{\Omega})s(\hat{\Omega})$ with $a(\hat{\Omega})$ as well as $a(\hat{\Omega})$ and $s(\hat{\Omega})^2$. In terms of the suboptimal estimators introduced before the correspond to $C_l^{a,s,a}$ and C_l^{s,a^2} respectively. In the second case analysis is exactly same as that of lensing reconstruction discussed before. However in the first case the optimal estimator is expressed as follows:

$$E_L^{s,aa}[\tilde{a}, \tilde{s}] = [N^{-1}]_{LL'} \{ Q_{L'}[\tilde{a}, \tilde{s}] - \tilde{a}_{lm} \langle \partial_{lm}^a Q_{L'}[\tilde{a}, \tilde{s}] \rangle - \tilde{s}_{lm} \langle \partial_{lm}^s Q_{L'}[\tilde{a}, \tilde{s}] \rangle \}. \quad (85)$$

The above estimator considering is same as Eq.(68); with the corresponding $Q_{L'}[\tilde{a}, \tilde{s}]$ function and its derivatives are given in Eq.(63).

The mixed bispectrum of CMB² – SZ is known to be exactly same as that of bispectrum we considered in the lensing reconstruction. This is true not only for SZ-lensing bispectrum but for other lensing-induced correlation-related bispectra too. The only difference is in different b_l s involved.

$$\begin{aligned} F_{LL'} = & 2\delta_{LL'} \sum_{ll'} \left\{ B_{LL' ll'}^{saa} B_{ll' ll'}^{saa} \frac{1}{C_L^{ss}} \frac{1}{C_l^{aa}} \frac{1}{C_l^{aa}} \right\} + \sum_l \left\{ 2 B_{LL' ll'}^{saa} B_{ll' ll'}^{saa} \left(\frac{C_L^{sa}}{C_L^{ss} C_L^{aa}} \right) \left(\frac{C_L^{sa}}{C_L^{ss} C_L^{aa}} \right) \frac{1}{C_l^{aa}} \right. \\ & \left. + B_{LL' ll'}^{saa} B_{ll' ll'}^{saa} \frac{1}{C_L^{ss}} \left(\frac{C_L^{sa}}{C_L^{ss} C_L^{aa}} \right) \left(\frac{C_l^{sa}}{C_l^{ss} C_l^{aa}} \right) + B_{LL' ll'}^{saa} B_{ll' ll'}^{saa} \frac{1}{C_l^{aa}} \left(\frac{C_L^{sa}}{C_L^{ss} C_L^{aa}} \right) \left(\frac{C_L^{sa}}{C_L^{ss} C_L^{aa}} \right) \right\}. \end{aligned} \quad (86)$$

This is an application of the case considered in Eq. (69). For the one-point mixed skewness associated with this power spectra the related Fisher error is simply given by the sum over all the elements. $F = \sum_{L,L'} F_{LL'}$. The cross correlations C_l^{sa} between the two maps $s(\hat{\Omega})$ and $a(\hat{\Omega})$ introduces the off-diagonal elements in the Fisher matrix even for the case of all-sky coverage and homogeneous noise. Ignoring the correlations we can recover the Fisher matrix elements derived for the case of lensing reconstruction.

In addition to considering the cross-correlation of $s(\hat{\Omega})$ and $a^2(\hat{\Omega})$ as discussed above, the other estimator of non-Gaussianity that we can consider is by considering the cross-correlation of product field $s(\hat{\Omega})a(\hat{\Omega})$ and $a(\hat{\Omega})$ which is same as the estimator is same as that defined in Eq. (74) with the relevant Q term and its derivative given by Eq. (73). The associated Fisher matrix is given by Eq. (76). The one-point estimator recovered from both of these degenerate estimators will be the same.

8 CONCLUSIONS

Extending previous work for estimation of power-spectra from correlated data sets we show how pseudo- C_l -based approaches (PCL) can be used for estimation of cross-correlation power spectra from multiple cosmological surveys through a joint analysis. Analytical results were derived under very general conditions using an arbitrary mask as well as arbitrary noise properties. We also keep the weighting of the data completely general. Our analytical results also include a systematic analysis of covariance of various deconvolved $\hat{C}_l$ s characterizing auto- and cross-power spectra from a joint analysis. While PCL-based approaches are known to be unbiased they are not in general optimal. However they can be made to act in a near-optimal way by the introduction of weights in different regimes corresponding to signal or noise domination. These studies will be useful in analyzing simulated as well as real survey data either in projection or in 3D. We specialise these expressions to recover well-known f_{sky} approximation used in the literature for the error analysis. Using a halo model inspired approach we compute the expected cross-correlation signal in cross-correlating NVSS type survey with the CMB sky through the ISW effect. We also study the cross-correlation between the frequency-cleaned SZ surveys against the ISW effect. The cross-correlation study also provides the covariances among different estimated C_l s and the signal-to-noise of detection for a specific survey. However

we want to stress that the formalism developed here is more powerful and can tackle many issues in analysing realistic surveys. A detailed study using simulations will be presented elsewhere.

The analysis of the bispectrum is one step beyond the power-spectrum and provides additional cosmological information. The primary motivation to date has been to put constraints on early-universe scenarios, however secondary bispectra can play a significant role in enhancing our understanding of large-scale structure formation scenarios. The secondary bispectrum is mainly related to mode-coupling by secondary effects and lensing. We study various statistics which can directly handle realistic data sets. Extending previous work by Munshi & Heavens (2009), we take into account multiple correlated fields which are used for constructing a mixed skewness at one-point level as well as constructing a skew spectrum at the level of two-point. A very general framework was developed for the study of bispectrum from correlated fields in an unbiased and optimized way. We introduce the inverse covariance weighting and specialize our results for the analysis of bispectrum originating from lensing-secondary correlations. A simple-minded approach which handles the noise and partial sky coverage in a nearly optimal way using Monte Carlo techniques is also discussed. We also develop an approach based on PCL to study the skew spectrum. This approach, whilst suboptimal, can handle the noise and partial sky coverage directly. It is also possible to use weights to make it near optimal in the limit of high l , and can be useful mainly because of its speed of handling MC realisations. In its most general form, the estimator E_L (equation (64) for the skew spectrum of mixed fields includes the effect of partial sky coverage and inhomogeneous noise, and provides a compact function which can be compared with theoretical models to identify the source of the correlations between the fields. The associated Fisher matrix (equation 66) allows statistical analysis of the E_L estimates, allowing the estimation of the relative contributions from different physical processes.

For specific examples we have focussed on probing the secondary non-Gaussianity with Planck type all-sky experiments and surveys such as NVSS. The signal to noise ratios for cross-correlation studies involving lensing potential and secondaries such as SZ and ISW would allow detection with Planck. However to differentiate among various effects one would need to go beyond cumulative signal-to-noise estimates and the statistics which we introduced here will be useful diagnostic tools.

There has been lot of work by a number of authors to detect correlations between the WMAP CMB and large scale structures, which typically conclude with a constraint on the dark energy (accelerating universe). Analysis of secondary bispectrum has also been attempted. However, at this point, consistent simulations which can correctly take into account, the correlation between CMB and the LSS, and the impact of the LSS on the various observables is still remains to be developed. Though a patchwork of simulations are getting ready, we still lack suitable simulations which can be used both for cross-correlational analysis or the entire range of bispectrum analysis at the moment. Our approach can be invaluable in quantifying accuracy of such consistency check and eventually to put constrain on cosmology using real high resolution data. We have not taken into account the errors or residuals from foreground removals. Some of the foreground contaminations may well be correlated to various LSS tracers. These issues and how PCL based approach can tackle them will be dealt with elsewhere.

9 ACKNOWLEDGEMENTS

DM was supported by a STFC rolling grant at Royal Observatory at Edinburgh, Institute for Astronomy, when this work was performed. It is a pleasure to acknowledge useful exchanges with Jacques Delabrouille, Matthias Bartelmann and Patricio Vielva Martinez. AC acknowledges support from NSF CAREER AST-0645427.

REFERENCES

- Acquaviva V., Bartolo N., Matarrese S., Riotto A., 2003, Nucl. Phys. B667, 119
- Alishahiha M., Silverstein E., Tong T., 2004, Phys. Rev. D70, 123505
- Arkani-Hamed N., Creminelli P., Mukohyama S., Zaldarriaga M., 2004, JCAP0404:001
- Afshrodi N., Loh, Y., Strauss, M.A., 2004, Phys. Rev. D 69, 083524
- Afshrodi N., 2004, Phys. Rev. D, 70, 083536
- Babich D., 2005, Phys. Rev. D72, 043003
- Babich D., Pierpaoli E., 2008, Phys. Rev. D77, 123011
- Babich D & Zaldarriaga M., 2004, Phys. Rev. D70, 083005
- Babich D., Creminelli P., Zaldarriaga M., 2004, JCAP, 8, 9
- Bartolo N., Matarrese S., Riotto A., 2006, JCAP, 06, 024
- Buchbinder E.I., Khoury J., Ovrut B.A., 2008, Phys.Rev.Lett,100:171302
- Boughn S., Crittenden R., MNRAS, 2005, 360, 1013
- Boughn S., Crittenden R., Nature 427 (2004) 45
- Boughn S., Crittenden R., New Astron.Rev. 49 (2005) 75-7
- Cabella P., Hansen F.K., Liguori M., Marinucci D., Matarrese S., Moscardini L., Vittorio N., 2006, MNRAS, 369, 819
- Cabre A., Gaztanaga E., Manera M., Fosalba P., Castander F. Mon.Not.Roy.Astron.Soc.Lett., 372 (2006) L23-L27
- Castro P., 2004, Phys. Rev. D67, 044039 (erratum D70, 049902)
- Chen X., Huang M., Kachru S., Shiu G., 2006, hep-th/0605045
- Chen X., Easter R., Lim E.A., 2007, JCAP, 0706:023
- Chen G. & Szapudi I., Astrophys.J, 647, 2006, L87-L90, 2006
- Cheung C., Creminelli P., Fitzpatrick A.L., Kaplan J., Senatore L., 2008, JHEP, 0803, 014
- Cooray A.R., Hu W., 2000, ApJ, 534, 533-550
- Cooray A., Hu W., Tegmark, M. 2000, ApJ, 540, 1-13

- Cooray A., 2001a, Phys. Rev. D, 64, 043516
 Cooray A., 2001b, Phys. Rev. D, 64, 063514
 Cooray A., 2006, Phys. Rev. Lett., 97, 261301
 Cooray A., Li C., Melchiorri A., 2008, Phys. Rev. D 77, 103506
 Cooray A., 2000, Phys. Rev. D, 62, 103506
 Cooray A. Seth R., Phys. Rept. 372 (2002) 1-129
 Creminelli P., 2003, JCAP 0310, 003
 Creminelli P., Nicolis A., Senatore L., Tegmark M., Zaldarriaga M., 2006, JCAP, 5, 4
 Creminelli P., Senatore L., Zaldarriaga M., Tegmark M., 2007, JCAP, 3, 5
 Creminelli P., Senatore L., Zaldarriaga M., 2007, JCAP, 3, 19
 Corasaniti, P.S., Giannantonio T., Melchiorri, A., 2005, Phys. Rev D 72, 023514
 Diego, J.M., Silk J., Sliwa W., New Astron.Rev. 47 (2003a) 855,
 Diego, J.M., Silk J., Sliwa W., Mon.Not.Roy.Astron.Soc. 346 (2003b) 940
 Efstathiou G., 2004, MNRAS, 349, 603
 Efstathiou G., 2006, MNRAS, 370, 343
 Falk T., Madden R., Olive K.A., Srednicki M., 1993, Phys. Lett. B 318, 354
 Fosabala, P. & Gaztanaga E., (2004), MNRAS, 350, L37
 Fosabala, P. & Gaztanaga E., Castander F., (2003), ApJ, 597, L89
 Gangui A., Lucchin F., Matarrese S., Mollerach S., 1994, ApJ, 430, 447
 Giannantonio, T. et al., Phys.Rev. D 74 (2006) 063520
 Goldberg D.M., Spergel D.N., 1999, Phys. Rev. D 59, 103002
 Gupta S., Berera A., Heavens A.F., Matarrese S., 2002, Phys.Rev. D 66, 043510
 Heavens A.F., 1998, MNRAS, 299, 805
 Hivon E., Górski K. M., Netterfield C. B., Crill B. P., Prunet S., Hansen F., 2002, ApJ, 567, 2
 Hirata C.M., Ho S., Padmanabhan N., Seljak U., Bahcall N., 2008, Phys.Rev.D 78, 043520
 Ho S., Hirata C. M., Padmanabhan N., Seljak U., Bahcall N., 2008, Phys.Rev.D 78, 043519
 Hu W., 2000, PhRvD, 62, 043007
 Hu W., Okamoto T., 2002, ApJ, 574, 566
 Jaffe A.H., Kamionkowski M., Phys.Rev. D 58 (1998) 043001
 Komatsu E., Spergel D. N., 2001, Phys. Rev. D 63, 3002
 Komatsu E., Spergel D. N., Wandelt B. D., 2005, ApJ, 634, 14
 Komatsu E., Wandelt B. D., Spergel D. N., Banday A. J., Górski K. M., 2002, ApJ, 566, 19
 Komatsu E., et al., 2003, ApJS, 148, 119
 Koyama K., Mizuno S., Vernizzi F., Wands D., 2007, JCAP 0711:024
 Linde A. D., Mukhanov V. F., (1997), Phys. Rev. D 56, 535
 Lyth D.H., Ungarelli C., Wands D., 2003, Phys. Rev. D 67, 023503
 Maldacena J.M., 2003, JHEP, 05, 013
 Medeiros J., Contaldi C.R., 2006, MNRAS, 367, 39
 Moss I., Xiong C., 2007, JCAP, 0704, 007
 Munshi D., Souradeep, T., Starobinsky, Alexei A., 1995, ApJ, 454, 552
 Munshi D., Heavens A., arXiv:0904.4478
 Munshi D., Melott A.L., Coles P., MNRAS, 2000, 311, 149.
 Nolte et al., Astrophys.J., 2004, 608, 10
 Padmanabhan N., Hirata C. M., Seljak U., Schlegel D., Brinkmann J., Schneider D.P. Phys.Rev. D 72 (2005) 043525
 Peiris H. V. & Spergel D.N., 2000, ApJ, 540, 605
 Sachs R. K. & Wolfe A.M., 1967, ApJ, 147, 73
 Salopek D. S., Bond J. R., 1990, PhRvD, 42, 3936
 Salopek D. S., Bond J. R., 1991, PhRvD, 43, 1005
 Santos M.G. et al., 2003, MNRAS, 341, 623
 Serra P., Cooray A., 2008, Phys. Rev. D, 77, 107305
 Smith K.M., Zahn O., Dore O., 2007, Phys. Rev. D, 76, 043510
 Smith K. M., Zaldarriaga M., 2006, arXiv:astro-ph/0612571
 Smith K.M., Senatore L., Zaldarriaga M., 2009, arXiv:0901.2572
 Spergel D.N., Goldberg D. M., 1999, Phys.Rev. D 59, 103001
 Spergel D.N. et al., 2007, ApJS, 170, 377
 Sunyaev R. A., Zeldovich I. B., 1980, MNRAS, 190, 413
 Szapudi I., Szalay A.S. Astrophys.J. 515 (1999) L43
 Tegmark M., Phys.Rev. D 55 (1997) 5895-5907
 Verde L., Spergel D.N., 2002, Phys. Rev. D 65, 043007
 Wang L., Kamionkowski M., 2001, Phys. Rev. D 61, 3504
 Yadav A. P. S., Wandelt B. D., 2008, PhRvL, 100, 181301
 Yadav A. P. S., Komatsu E., Wandelt B. D., Liguori M., Hansen F. K., Matarrese S., 2008, ApJ, 678, 578

18 *Munshi, Valageas, Cooray & Heavens*

Yadav A. P. S., Komatsu E., Wandelt B. D., 2007, ApJ, 664, 680

Zahn O., Zaldarriaga M., 2006, ApJ, 653, 922